

SCREENING YEAST STRAINS FOR ALCOHOL FERMENTATION FROM THE DRIED TRADITIONAL YEAST

PHAM NGUYEN PHUONG THAO, BUI THI THANH THUY

University of Science, Vietnam National University Ho Chi Minh City - pnpthao@hcmus.edu.vn

LE BUI TRUNG TRINH, NGUYEN MY PHI LONG

University of Science, Vietnam National University Ho Chi Minh City

(Received: February 06, 2018; Revised: March 30, 2018; Accepted: April 16, 2018)

ABSTRACT

Methods to produce rice alcohol by dried traditional yeast are unstable because the yeast system in dried traditional yeast has depended on nature and not been controlled. In this study, a total of 15 different kinds of dried traditional yeast were prepared and screened. Each yeast strain was evaluated by analyzing its fermentation property and alcohol tolerance. There are 19 yeast strains were isolated and their growth conditions and ethanol producing properties were examined. Results showed that three strains S1, BT, BL3 grew and produced ethanol at temperature 28-30°C, and pH 5-5.5. Especially, high concentration ethanol tolerance ability of the three strains was at 8-18%. Our results showed that these strains were valuable microorganisms and could be utilized as a basis for further study of dried traditional yeast in traditional alcoholic beverages.

Keywords: Rice alcohol; The dried traditional yeast; Yeast.

1. Introduction

As the society growth there is a strong demand for a variety of delicious and national representative wines. Moreover, it has to be ensured that these wine is safe to the users. If Japan is proud of its Sake (Iwata, H., Suzuki, 2003), Korea has Sochou or Western European countries has Vodka (Russia) and Whiskey (Scotland), rice alcohol is considered as an alcoholic beverage commonly used in Vietnam basing on its wet rice civilization. There is no accurate record of how Vietnam's traditional alcohol was made. However, it has been known from the ancient times that rice alcohol has been used extensively in holidays, festivals, sacrifices and even in regular meals.

Alcohol in Viet Nam is produced in many different ways. Each area in Viet Nam has a distinct taste of alcohol such as Mau Son - Lang Son, Dao - Yen Bai, Bau Da - Binh Dinh, Go Den - Long An. Traditional method to produce alcohol using dried yeast is still

widely applied, although the efficiency and the product in many regions are still unstable. This instability is due to the microorganisms present in the dried yeast without being controlled. They play an important role in transforming substances from raw materials into sugars and then to liquors through saccharification process and fermentation. They make a significant influence on food quality parameters such as taste, texture and nutritious value (Kofi et al., 2006). There are various types of yeast involved in the production of alcohol such as *Hanseniaspora*, *Candida*, *Metschnikowia*... (Ray, B., 2001; Schutz M. and Gafner J., 1993). Especially, *Saccharomyces* strains are used majority in many fermentation process (Dung et al., 2007).

Around the world, there were several research on selecting high activity yeast strains for alcohol fermentation. A study of F. Noé Arroyo-López et al. done in 2009 examined the

effects of heat, pH and concentration of sugar to 3 strains of *S. cerevisiae* T73, *S. kudriavzevii* IFO 1802T and hybrid strain *S. cerevisiae* × *S. kudriavzevii* W27 used in wine making. In particular, the W27 hybrid strain has the ability to ferment into alcohol at high pH (5.5) and low sugar concentration (10g/l). T73 and W27 could be used in industrial production. In 2011, H. Yamamoto isolated 7 yeast strains with high alcohol fermentation activity and Shochu flavor. Particularly, the MF062 strain had highly fermentable activity, high yield, incense and heat tolerance. The results of genetic and phylogenetic datashow that MF062 is *Saccharomyces cerevisiae* but has many different characteristics from industrial yeasts using in shochu production. In this year, Yeon-Ju Lee¹ et al. surveyed the ability of alcohol and osmotic pressure tolerance of different yeast strains using maltose and hydrolysis of starch. In the 637 presumed yeast species, 115 strains showed good growth in yeast-peptone-dextrose containing 30% dextrose, 7% alcohol, or 2% maltose, and produced 5 α -amylase enzymes. Analyzing the nucleotide sequence of the 26S rDNA gene classified these yeast strains into 13 species of five genera: *Pichia anomala* was the most common (41.7%), *Wickerhamomyces* (19.2%), *P. guilliermondii* (15%), *Candida* spp. (5.8%), *Kodamaea ohmeri* (2.5%) and *Metschnikowia* spp. (2.5%). The isolation NK28, identified as *Saccharomyces cerevisiae*, had all the desired properties for the purposes of this study, except for the production of α -amylase but the ability to ferment the wine better than commercial wine yeast counterpart. Another study, conducted in 2015, evaluated the fermentation efficiency in producing vodka from potato of three strains *Saccharomyces cerevisiae*. Results showed that bread yeast was most effective in the medium supplemented with hydrolysed potatoes and sucrose (17% by weight):

substituting substrates into high yields (each gram sugar lost, 0.47g ethanol created), high fermentation product (91.4%), high ethanol content (6.05%/V).

In Viet Nam, there were many reports about yeast for producing alcohol isolated from different sources of raw materials like starch from rice. In 2009, Ngo Thi Phuong Dung surveyed fermentation and alcoholic tolerance of different yeast strains. From 50 germs isolated from wine yeast, 9 yeast lines with fast and strong fermentation in glucose solution after 14 hours of fermentation were selected. 9 broths of yeast collected from fermentation in reducing sugar solution were able to use and almost completely alter sugar reduction at 18% (w/v). Also, these yeast could produce ethanol with concentration of 8.3 – 8.6% (w/v). The project also determined the alcoholic beverage of yeast strains and found that these seven yeast strains of *Saccharomyces* belonged to *Ascomycetes*. Another research on the production of some traditional wine from rice had been done by engineer Nguyen Dinh Quy from 2010-2012. The results indicated that currently locally available microorganisms contained three main groups: mold, yeast, and imitation yeast. The author selected four strains suitable for yeast production. The research has developed a technological process to manufacture enzyme with highly enzymatic, alpha amylase activity at 1102 U/g and glucoamylase at of 241 U/g. Also in 2012, the research team led by Nguyen Huu Thanh et al. conducted a number of studies on biological characteristics and identified yeast which were isolated from the dry traditional yeast in the Cuu Long Delta. 128 strains were isolated with 30 strains tolerate temperature at 50°C and alcohol tolerance of 17 ml/l. Sequencing was done in 10 strains, 7 strains were identified as *Saccharomyces cerevisiae*, 3 strains were *Clavispora lusitaniae*. Complete the technology and equipment of traditional rice

wine production with 800,000 l/year. To be conducted in 2013-2015 by Dr. Nguyen Viet Anh as the director. The project had completed the process of technology production of industrial rice wine which increased from 300,000 liters per year to 800,000 liters per year. Rice wine product reached good quality, stable, aromatic similar to the traditional rice wine, no toxic impurities, harmful. Designing and manufacturing mold production equipment scale of 1,500 kg/batch replaced manual mold culture, passive, easy to contaminate. In addition, the project has developed the technological process and production model of fermented rice vinegar by submerged fermentation using waste by-products of the process of producing industrial rice wine. The quality of the vinegar produced was much higher (8%) than the surface fermentation, characteristic aroma, short fermentation time. The project was accepted on 24.6.2016 with excellent results.

Therefore, the aim of this study was to select yeast strains with high alcohol-fermented ability and investigate optimum conditions for alcohol fermentation.

2. Materials and methods

2.1. Isolation and identification of yeast strains

The fifteen different dried traditional yeast were collected from Long An market, Go Den market and Cho Lon market → enriched in liquid Hansen medium (pepton 10 g/l, MgSO_4 3 g/l, KH_2SO_4 3 g/l, glucose 50 g/l) (Use a scalpel knife to remove the powder in the middle of the dried traditional yeast) incubated for 24 hrs at 28-30°C → Isolated by streaking on Hansen agar plates → selecting individual characteristic colonies to purify by streaking on Hansen agar and stored.

The yeast strains were identified to genus based on their morphological characteristics and the bud forming and spore forming with the classification key of Kurtzman and Fell,

1998 (Kurtzman, C.P. & Fell, J.W., 1998).

All isolated yeast strains were stored in Hansen medium containing 40% glycerol at -30°C.

2.2. Investigation of fermentation property of yeast strains

The fermentation test of 19 strains was conducted in a molasses medium with a concentration of 15% sucrose. 30 ml yeast biomass (10^8 cells/ml) was added in 300 ml molasse medium, incubated at 20°C. The speed of CO_2 generation of strains was determined through the volume of CO_2 produced in the falcon tube upside down by the time. The yeast strain has a faster fermentation capacity is the yeast strain that produces higher volume columns over time




Figure 1. System diagram used to determine the speed of CO_2 generated in Fermentation properties test

2.3. Investigation of ethanol tolerance of yeast strains

Test tubes containing 5 ml of liquid Hansen medium were sterilized at 121°C for 15 minutes. Ethanol was added to the medium with corresponding concentrations of 8%, 10%, 12%, 14%, 16%, 18%, 20%. 1 ml biomass (10^8 cells/ml) of 19 strains were inoculated into 19 test tubes, incubated at room conditions. After 24 hrs, the effervescence in test tubes was recorded and this solution of yeast was streaked on Hansen agar plate to verify for effervescence is due to the fermentation of yeast through ability to form colonies. The yeast strains produced gas in high alcohol medium and form colonies on agar plate were the ethanol tolerance strains.

2.4. Investigation of optimum conditions for alcohol fermentation

To survey of optimum initial pH for the fermentation, 10% volume yeast (10^7 cells/ml) was inoculated in each erlen containing 200 ml liquid Hansen medium with different pH values of 4.0, 4.5, 5.0, 5.5 and 6.0; incubated at 28°C for 24 hrs.

Optimal sucrose concentration was determined by adding 10% volume yeast solutions (10^7 cells/ml) in erlens containing 200 ml liquid Hansen medium with different sucrose concentrations of 5%, 7%, 9%, 11%, 13%, 15%, 17%, 19% and 21%; incubated at 28°C for 120 hrs.

To study the effect of temperature on the fermentation, 10% volume yeast (10^7 cells/ml) was inoculated erlens with 200 ml liquid Hansen medium, incubated for 24 hrs at different temperatures of 26°C, 28°C, 30°C and 32°C.

After the incubation time, alcohol in

experiment samples was distilled by fractional distillation and alcohol concentration was determined by alcoholmeter. For experiment to determining the suitable sucrose concentration for fermentation, the total remaining sugar was evaluated after fermentation by Phenol-Sulphuric acid method (Nielsen, S. S., 2010). All tests are repeated 2 times independently and in each time, a sample is made. The result is the average value of two times the experiment and Standard deviation.

3. Results and Discussion

3.1. Results of isolation and identification


Based on morphological characteristics (Table 1, Fig. 2, Fig. 3), 19 isolated strains belong to genus *Saccharomyces*. They had multilateral budding, produced little or no acetic acid. Ascospores were formed and asci were globose. Gelatin media was not strongly liquefied, and the strains was not assimilate nitrate.

Table 1


Characteristics of cells and colonies of cultured yeast in liquid hansen media

STT	Strain code	Cell shape	Cell size (µm)	Colony shape
1	AD	Small oblong	2,0 x 5,0	yellowish white, dry, rounded margin, high rise, smell of alcohol, size 3 mm.
2	S1	Oblong	1,5 x 5,5	yellowish white, rounded margin, high rise, smell of alcohol, size 2-3 mm.
3	TN	Oval	1,5 x 3,5	Ivory white, dry, rounded margin, high rise, size 2-3 mm
4	HG	Obovoid	4,5 x 5,0	Ivory white, rounded margin, large size, slightly mucus, high rise, small of alcohol, size 2-5 mm.
5	TH	Obovoid	3,0 x 5,0	Slightly yellow, rounded, small size, high rise, wrinkle margin, size 3 mm.
6	BC1.1	Oval	2,0 x 4,0	Large size, powdered, dry, high rise, wrinkle margin, centered in the middle, odourless, size 5mm.
7	BC1.2	Oval	1,5 x 3,5	White, large size, powdered, high rise, wrinkle margin, Odoriferous, size 4-5 mm
8	VL	Oval	3,5 x 5,5	White, rounded margin, high rise, slightly mucus, odourless, size 2-3 mm
9	BL1	Oval	4,5 x 5,0	White, high rise, wrinkle margin, centered in the middle, mucus, odoriferous, size 5-6 mm.
10	BL2	Oblong	2,0 x 5,0	White, high rise, small size, rounded margin, mucus, odoriferous, size 2-3 mm


STT	Strain code	Cell shape	Cell size (µm)	Colony shape
11	BL3	Spherical	2,0 x 3,5	Milky, high rise, wrinkle margin, dry, odoriferous, size 3-4 mm.
12	BC2.1	Oval	1,5 x 3,5	White, small size, high rise, wrinkle margin, dry, odourless, size 2-3 mm.
13	BC2.2	Oval	2,5 x 4,5	Milky, wrinkle margin, centered in the middle, dry, odourless, size 2-3 mm.
14	BC2.3	Oblong	2,0 x 6,0	White, centered in the middle, dry, The margin has roots, size 5-6 mm.
15	MB	Oval	2,5 x 5,0	Slightly milky, rounded margin, mucus, smell of alcohol, size 3-5 mm.
16	TH	Oblong	2,0 x 4,0	Yellowish white, small size, high rise, wrinkle margin, smell of alcohol, size 3 mm
17	BC3	Oblong	2,5 x 4,5	White, small size, high rise, mucus, centered in the middle, odourless, size 2-3 mm.
18	HN	Small oval	3,0 x 3,5	Yellowish white, large size, rounded margin, high rise, mucus, smell of alcohol, size 5 mm.
19	BT	Spherical	4,5 x 5,5	Slightly milky, centered in the middle, rounded margin, dry, smell of alcohol, size 5 mm.
20	TX	Spherical	2,5 x 5,0	Milky white, high rise, small size, dry, rounded margin, odourless, size 2-3 mm.




BC23: Oblong




BL1: Oval




BT: Spherical



HG: Obovoid



HN: Small oval



AD: Small oblong

Figure 2. Types of morphological characteristics of isolated strains in liquid hansen media.
Scale bar 5 µm.




Figure 3. Location of yeast bud scars in (arrow) liquid hansen media for 20 days. Scale bar 5 μ m.




Figure 4. Morphological characteristics of asci with 4 ascospores of the yeast strain BC2.2 (arrow). Scale bar 10 μ m.




Figure 5. A: Gelatin medium was not strongly liquefied by growth of yeast strains. B: compared to the control sample (+), the S1 strain formed turbid solution in the tube but gelatin was not liquefied at 20°C, C: the strain S1 did not grow on Bacto-Yeast Carbon base media added Nitrate.

Note: The control sample (+) is Gelatin medium.

3.2. Fermentation property and ethanol tolerance of yeast strains

The starting time of fermentation is different between 19 yeast strains even they

were inoculated in the same condition. Among them, 3 strains S1, BL3, BT had the earliest fermentation time 83 mins, 89 mins và 99 mins, respectively (Fig. 6).




Figure 6. The CO₂ generation speed of yeast strains

According to Brown et al (1981) and Sa-Correia et al (1983), ethanol inhibits both growth and fermentation of yeasts (Kunkee R. E. and Bisson L. F., 1993; Reed G. and Nagodawithana T. W., 1991). This study determined ethanol tolerance of the isolated strains by producing biomass and CO₂ gas in the test tube (Table 2), then they were streaked on agar plates for verification

(Fig. 7). Results showed that the biomass of yeast decreases as the concentration of ethanol in the solution increases. At concentrations of ethanol 8% to 14%, the biomass obtained from 19 strains. At a concentration of ethanol 18% only a few yeast strains grew. Particularly with 20% ethanol concentration, no strains survived. Ethanol might inhibit metabolic capacity of yeast.

Table 2

Alcohol tolerance of yeast strains

Strain	Alcohol content (%)						
	8	10	12	14	16	18	20
BT	+	+	+	+	+	-	-
TN	+	+	+	+	-	-	-
BC2.3	+	+	+	+	-	-	-
AD	+	+	+	+	+	+	-
HN	+	+	+	+	+	+	-
BC1.1	+	+	+	+	+	-	-
BC3	+	+	+	+	+	+	-
S1	+	+	+	+	+	+	-
BL2	+	+	+	+	-	-	-

Strain	Alcohol content (%)						
	8	10	12	14	16	18	20
VL	+	+	+	+	+	-	-
BL3	+	+	+	+	+	-	-
BC2.1	+	+	+	+	+	-	-
BC2.2	+	+	+	+	-	-	-
HG	+	+	+	+	+	-	-
BL1	+	+	+	+	-	-	-
MB	+	+	+	+	-	-	-
TX	+	+	+	+	+	-	-
TH	+	+	+	+	-	-	-
BC1.2	+	+	+	+	-	-	-

Note: +: biomass and CO₂, -: non biomass and CO₂




Figure 7. The alcohol tolerance of yeast. (A) CO₂ produced in test tube, (B) the growth of yeast biomass on agar plate for verification.

3.3. Optimum conditions for alcohol fermentation

Based on the fermentation property and high ethanol tolerance, three strains S1, BT, BL3 were chosen in the next experiments.

3.3.1. Effect of initial pH on alcohol fermentation

This study evaluated the effect of initial pH on the fermentation. The optimum pH was determined by measuring the alcohol concentration after 24hrs incubated at 28°C. The highest concentration of alcohol was found at pH 5 to 5.5, depending on the strain (Fig. 8). At pH of 4 or 6 the alcohol concentration was lower.




Figure 8. Effect of pH on the production of alcohol of three strains S1, BL3 and BT in 24 hrs.

3.3.2. Effect of temperature on alcohol fermentation

Temperature effected ethanol yields, by-products of the fermentation, and aroma of the wine. At the higher temperatures, the yield of ethanol is slightly decreased. Conversely, at lower temperature, yeast grows slowly. The rate of fermentation increases with increasing

temperature up to about 30° to 33°C (Reed G. and Nagodawithana T. W., 1991). This study result showed that the suitable temperature for the fermentation of the strains is from 28 to 30°C and at 30°C the highest yield of alcohol production (Fig. 9). According to Luong Duc Pham (2006) this is also the suitable temperature for the development of biomass.




Figure 9. Effect of temperature on the production of alcohol of three strains S1, BL3 and BT for 24 hrs.

3.3.3. Effect of sucrose concentration on alcohol fermentation

According to Reed (1991), the species *S. cerevisiae* is capable of using sucrose for growing. In this study, most isolates belong to the *Saccharomyces* genus and sucrose was

used as the sugar source.

Each yeast strain needed different sucrose concentration to the maximum alcohol production. The strain S1 produced the highest alcohol concentration 7.8% in medium containing 19% sucrose (Fig. 10). Because

medium has low ethanol concentrations, the growth of the yeast is inhibited and little growth occurs above 9% ethanol (Reed G. and Nagodawithana T. W., 1991). In ethanol tolerance test, yeast cells could exist in medium with high ethanol concentration (19%) but their metabolism might be inhibited, resulting in reduced fermentation.

In table 3, the concentration of sucrose remaining after fermentation were nearly similar between there strains in all experimented sucrose concentrations, but alcohol concentration was decreased from 17% or 19% sucrose (Fig.10). It means that the fermentation process is inhibited when sucrose reaches 21%.




Figure 10. Effect of sucrose concentration to ethanol production of 3 strains S1, BL3 and BT in 5 days

Table 3

The sucrose remained after fermentation

Strain	Sucrose concentration (%)								
	5	7	9	11	13	15	17	19	21
S1	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8	3.8
BL3	3.8	3.7	3.7	3.8	3.7	3.7	3.8	3.7	3.8
BT	3.7	3.8	3.7	3.7	3.7	3.7	3.7	3.7	3.8

4. Conclusion

In this study, 19 yeast strains belong to genus *Saccharomyces* were isolated. There were 4 strains AD, HN, BC3, and S1 survived in medium with 18% ethanol. The concentration of sucrose remaining after their fermentation was from 3.77 to 3.80%. The

strain S1 can survive in medium added highest ethanol 18% and ferment in medium containing 19% sucrose and generated alcohol was 7.8% after 5 days cultured. This study could contribute preliminary data for further study of dried traditional yeast in traditional alcoholic beverages and their valuable microorganisms■

References

- Dung, N.T.P., Rombouts, F.M., and Nout, M.J.R. (2007). Characteristics of some traditional Vietnamese starch-based rice wine fermentation starters. *Food Science and Technology*, 40(1), 130-135.
- Iwata, H., Suzuki (2003). Purification and Characterization of Rice α -Glucosidase, a key Enzyme for Alcohol Fermentation of Rice polish. *Journal of Bioscience and Bioengineering*, 95(1), 106-108.
- Kofi, E. A., Nout, M. J. R., Sarkar, P. K. (2006). Occurrence and function of yeasts in Asian indigenous fermented foods. *FEMS Yeast Res*, 6(1), 30-39.
- Kunkee R. E. and Bisson L. F. (1993). *Wine making yeasts*. In: *The Yeast*. 2nd edn (AH Rose and JS Harrison, eds). Academic Press. New York.
- Kurtzman, C.P. & Fell, J.W. (1998). *The yeast, A taxonomic study*. Elsevier. Amsterdam.
- Luong Duc Pham (2009). *Nam men Cong Nghiep*. 2nd edn. Hà Nội: Khoa học và Kỹ thuật.
- Nielsen, S. S. (2010). *Phenol-Sulfuric Acid Method for Total Carbohydrates*. In: S.S. Nielsen. *Food analysis laboratory manual*. Food Science Texts Series.
- Ray, B. (2001). *Microorganism used in food fermentation*. In fundamental food microbiology.
- Reed G. and Nagodawithana T. W. (1991). *Yeast technology*. 2nd ed. Van Nostrand Reinhold Co., Inc., New York.
- Schutz M. and Gafner J. (1993). Analysis of yeast diversity during spontaneous and induced alcoholic fermentations. *Journal of Applied Bacteriology*, 75, 551-458.