



Research Article

MONOARYLATION OF DIPHENYLDITHIENYLETHENE BY LIGAND-FREE PALLADIUM CATALYZED C-H ACTIVATION

Le Tin Thanh^{1*}, *Nguyen Thi Phuong Nhi*¹, *Nguyen Hien*²,
*Le Thi Hong Hai*², *Le Thanh Thanh*³, *Dang Thanh Tung*⁴

¹Ho Chi Minh City University of Education, Ho Chi Minh City, Vietnam

²Ha Noi National University of Education, Ha Noi, Vietnam

³Faculty of Petroleum, PetroVietnam University, Ba Ria-Vung Tau Province, Vietnam

⁴Laboratoire de Physicochimie des Polymères et des Interfaces, Universités de CY Cergy Paris

*Corresponding author: *Le Tin Thanh* – Email: thanhlth@hcmue.edu.vn

Received: May 25, 2021; Revised: June 20, 2021; Accepted: September 01, 2021

ABSTRACT

This work reports on a green and convenient methodology for the arylation of diphenyldithienylethene by direct ligand-free palladium-catalyzed C–H bond functionalization. Based on this approach, a new diphenyldithienylethene-based compound with aryl substituent **DTE-3-CHO-1** was successfully synthesized from thiophene via 4 steps. These compounds were structurally identified by NMR and HR-MS spectral analyses.

Keywords: aggregation-induced emission; Diphenyldithienylethene; direct palladium-catalyzed arylation; fluorescence

MONO ARYL HOÁ DIPHENYLDITHIENYLETHENE THÔNG QUA PHẢN ỨNG HOẠT HOÁ TRỰC TIẾP LIÊN KẾT C-H SỬ DỤNG XÚC TÁC PALLADIUM KHÔNG LIGAND

¹Trường Đại học Sư phạm Thành phố Hồ Chí Minh, Thành phố Hồ Chí Minh, Việt Nam

*Tác giả liên hệ: *Lê Tín Thanh* – Email: thanhlth@hcmue.edu.vn

Ngày nhận bài: 25-5-2021; ngày nhận bài sửa: 20-6-2021; ngày duyệt đăng: 01-9-2021

TÓM TẮT

Bài báo trình bày một phương pháp hiệu quả để thực hiện aryl hoá diphenyldithienylethene thông qua phản ứng hoạt hoá trực tiếp liên kết C-H sử dụng xúc tác palladium không ligand. Hợp chất diphenyldithienylethene **DTE-3-CHO-1** được tổng hợp từ thiophene qua bốn giai đoạn. Giai đoạn chìa khoá của quá trình tổng hợp **DTE-3-CHO-1** là phản ứng hoạt hoá trực tiếp liên kết C-H được thực hiện giữa diphenyldithienylethene **3** và 3-bromobenzaldehyde sử dụng xúc tác Pd(OAc)₂, base KOAc trong dung môi DMAc tại 120 °C trong 16h. **DTE-3-CHO-1** thu được với

Cite this article as: Le Tin Thanh, Nguyen Thi Phuong Nhi, Nguyen Hien, Le Thi Hong Hai, Le Thanh Thanh, & Dang Thanh Tung (2021). *Using creative methodology to explore factors influencing teacher educator identity. Ho Chi Minh City University of Education Journal of Science, 18(9), 1359-1367.*

hiệu suất 49%. Hợp chất **DTE-3-CHO-1** tổng hợp được hứa hẹn là một hợp chất có các tính chất quang lý hấp dẫn. Cấu trúc của diphenyldithienylethene **DTE-3-CHO-1** cũng như các hợp chất trung gian trong quá trình tổng hợp được xác định dựa vào phổ NMR và HR-MS. Hợp chất **DTE-3-CHO-1** là một hợp chất mới.

Từ khóa: aggregation-induced emission; Diphenyldithienylethene, phản ứng aryl hoá trực tiếp xúc tác palladium, fluorescence

1. Introduction

Over the past decades, luminescent organic materials have been developed and applied into many research fields including photoelectronic devices, fluorescent sensors, bioimaging trackers, and photoluminescence therapy. In particular, aggregation induced emissions (AIE) have gained increasing attention on account of their fundamental importance and high promising potential applications. There are many compounds showing the AIE effects such as tetraarylethenes, multi-substituted alkenylated benzenes, heteroatom-bridged pentacyclic compounds (1,1-disubstituted tetraphenylgermole, *p*-aryltetraphenylphospholes, 1,1-diphenyltetraphenylsiloles, tetrabromothiophene, *N*-phenyltetraphenylpyrrole, tellurophenes, 1,1-disubstituted tetraphenylstannoles), and boron diketonates (Mei et al., 2015). Among them, tetraphenylethene has been widely developed due to its facile synthesis and straightforward post-functionalization for multi-purposed applications (Hong et al., 2011). The synthesis of tetraarylethenes were based on titan-assisted McMurry coupling of ketones. Normally, this approach gave low yields with many accompanying isomer products. Recently, there have been two efficient methodologies to construct multiarylated ethenes with high yields based either on , palladium-catalyzed *syn*-direct diarylation of aryldiethynyl *N*-methyliminodiacetyl boronates (Lin et al., 2019) or palladium catalyzed consequential multi-Suzuki reaction of di-, tri- and tetraboronated ethenes (Zhang et al., 2020).

Post-monofunctionalized TPEs have applications in various research fields thank to their interesting physical properties (Li et al., 2013). For example, 2-formylthienyl-TPE showed bright green color mechanoluminescence upon pressing or grinding under daylight at room temperature (Rananaware et al., 2015). 2-PyrylTPE was observed to quench light emissions in solution by π - π stacking of luminophors and to exhibit the AIE characteristics in the condensed phase by the restriction of intramolecular rotation (RIR) (Zhao et al., 2011). TPE carboxylic acid has been studied for its application in photobleaching and bioimaging activity towards bacteria including several species of gram-negative and gram-positive (Shi et al., 2019; Liu et al., 2017). Almost all of the post-monofunctionalized TPEs were synthesized after two steps, that are bromination and the classical palladium catalyzed coupling reactions or the metal halide exchanged reaction (Zhang et al., 2014).

To give a convenient approach for post-monofunctionalisation of diphenyldithienylethenes (DTE) that have dual fluorescent and aggregation induced

emission effects, we have studied a synthetic protocol to synthesize monoarylated DTE by a direct ligand-free palladium-catalyzed arylation of DTE with 3-bromobenzaldehyde.

2. Materials and methods

2.1. Instrumentation

NMR spectroscopic data were acquired on a Bruker Avance III at 500 MHz for ^1H -NMR and 125 MHz for ^{13}C -NMR. HR-MS spectra were recorded on a Bruker MICROTOF-Q 10,187.

2.2. Material

Reagents and solvents were obtained from commercial suppliers and were used without further purification. Column chromatography was carried out using Merck Kieselgel 60 silica gel (particle size: 32-63 Å). Analytical TLC was performed using Merck precoated silica gel 60 F-254 sheets. All the CH activation reactions were carried out under nitrogen atmosphere.

2.3. Synthesis

Di(thiophen-2-yl)methanone 1: A mixture of 2-thienoic acid (1.2 g, 0.01 mol, 1 eq), thiophene (0.84 g, 0.01 mol, 1 eq), and phosphorus pentoxide (1.6 g, 0.12 mol, 1.2 eq) in anhydrous toluene (10 mL) was refluxed for 2h. Another portion of phosphorus pentoxide (1.6 g, 0.12 mol, 1.2 eq) was added and the mixture was continued to reflux for another 2h. The precipitate was filtered off and the combined filtrates were washed with water (3 x 50 mL), 1M aq. NaOH (3 x 50 mL), dried over Na_2SO_4 , concentrated *in vacuo* to give a residue which was chromatographed over silica gel (*n*-hexane: ethyl acetate (5:1)) to give **1** (970 mg, 50 %) as a white solide. ^1H -NMR δ_{H} (500 MHz, CDCl_3 , δ ppm): δ 7.91 (2H, *dd*, J = 4.0 Hz, J = 1.0 Hz), δ 7.70 (2H, *dd*, J = 5.0 Hz, J = 1.0 Hz), δ 7.19 (2H, *dd*, J = 5.0 Hz, J = 4.0 Hz); ^{13}C -NMR δ_{C} (125 MHz, CDCl_3 , δ ppm): δ 178.8, 142.9, 133.4, 133.1, 127.9 (Bottalico et al., 2009).

1,1-Bis(thiophen-2-yl)-2,2-dibromoethene 2: **1** (194 mg, 1 mmol, 1.0 eq), CBr_4 (662 mg, 2 mmol, 2.0 eq) and PPh_3 (1.05 g, 4 mmol, 4.0 eq) were dissolved in dry toluene (8.0 mL). The solution was refluxed for 24h. After filtration, the filtrate was washed with water (3 x 50 mL), dried over Na_2SO_4 and concentrated *in vacuo*. Chromatography on silica gel (*n*-hexane) afforded **2** (150 mg, 43 %) as a red oil. ^1H -NMR δ_{H} (500 MHz, CDCl_3 , δ ppm): δ 7.39 (2H, *dd*, J = 5.0 Hz, J = 1.0 Hz), δ 7.10 (2H, *dd*, J = 4.0 Hz, J = 1.0 Hz), δ 7.00 (2H, *dd*, J = 5.0 Hz, J = 4.0 Hz) (Chang et al., 2016).


1,1-Bis(thiophen-2-yl)-2,2-diphenylethene 3: A mixture of **2** (70 mg, 0.2 mmol, 1.0 eq), PPh_3 (26 mg, 0.1 mmol, 0.5 eq), Na_2CO_3 (106 mg, 1.0 mmol, 5.0 eq), $\text{Pd}(\text{OAc})_2$ (5 mg, 0.022 mmol, 0.1 eq) and phenylboronic acid (122 mg, 1.0 mmol, 5.0 eq) in dioxane : H_2O 4:1 (v:v) (10 mL) was refluxed overnight. After the completion of the reaction, the precipitate was filtered and the filtrate was washed with water (3 x 50 mL). The organic layer was dried over Na_2SO_4 and concentrated *in vacuo*. The residue was purified by

column chromatography on silica gel (*n*-hexane) to yield **3** (38 mg, 55 %) as a white solid. $^1\text{H-NMR}$ δ_{H} (500 MHz, CDCl_3 , δ ppm): δ 7.19-7.13 (12H, *m*), 6.80 (2H, *dd*, $J = 5.0$ Hz, $J = 4.0$ Hz), 6.76 (2H, *dd*, $J = 4.0$ Hz, $J = 1.0$ Hz) (Wang et al. 2018); $^{13}\text{C-NMR}$ δ_{C} (125 MHz, CDCl_3 , δ ppm): δ 145.6, 143.2, 142.3, 130.6, 129.7, 128.0, 127.0, 126.6, 126.3, 126.1.

3-(5-(2,2-Diphenyl-1-(thiophen-2-yl)vinyl)thiophen-2-yl)benzaldehyde DTE-3-CHO-1: A suspension of **3** (69 mg, 0.2 mmol, 1.0 eq), $\text{Pd}(\text{OAc})_2$ (1.0 mg, 0.006 mmol, 0.03 eq), KOAc (58 mg, 0.6 mmol, 3.0 eq) and 3-bromobenzaldehyde (3.0 eq) in DMAc (5.0 mL) was stirred at 120 °C for 16h. All the insolubles were filtered and the combined filtrates were washed with water (3 x 30 mL), dried over Na_2SO_4 , and the solvent was evaporated *in vacuo* to give a crude residue which was purified by column chromatography on silica gel (*n*-hexane:ethyl acetate 9:1, v/v) to give the pure monoarylated product **DTE-3-CHO-1** (44 mg, 49%) as a yellow solid. $^1\text{H-NMR}$ δ_{H} (500 MHz, CDCl_3) δ 10.00 (s, 1H), 7.96 (t, $J = 2.0$ Hz, 1H), 7.74 – 7.69 (*m*, 2H), 7.48 (t, $J = 7.5$ Hz, 1H), 7.22 – 7.14 (*m*, 11H), 7.11 (*d*, $J = 4.0$ Hz, 1H), 6.87 – 6.81 (*m*, 2H), 6.73 (*d*, $J = 4.0$ Hz, 1H). $^{13}\text{C-NMR}$ δ_{C} (125 MHz, CDCl_3) δ 192.0, 145.2, 143.1, 143.0, 142.9, 142.8, 136.9, 131.1, 130.9, 130.6, 130.6, 129.9, 129.5, 128.4, 128.2, 128.0, 127.3, 127.1, 126.6, 126.4, 126.3, 123.3. HR-ESI-MS: calcd $[\text{C}_{29}\text{H}_{21}\text{OS}_2]^+$ ($[\text{M}+\text{H}]^+$) = 449.1034, found = 449.1012.

3. Results and discussion

Diphenyldithienylethene was synthesized by two methodologies. First, dissymmetric 1,1-diphenyl-2,2-(hetero-)arylethene was synthesized by Suzuki and/or Stille coupling reaction of corresponding gem-dibromoethene and boronic or stannyl (hetero-)aryl (Zhang et al., 2014). One year later, diphenyldithienylethene was synthesized by desulfurization of thiiranes (Młostoń et al., 2016). This method was known as ‘two-fold extrusion reaction’ and needed further two steps for synthesis of starting materials including diaryldiazomethane and dithienyl thioketones or dithienyldiazomethane and diaryl thioketones. In our strategy, we have modified the synthesis of diphenyldithienylethene by reversed both reactants via gem-dibromodithienylethene and phenylboronic acids. Gem-dibromodithienylethene was synthesized after two steps as described in Scheme 1. The Suzuki reaction was employed for construction of tetrasubstituted ethene **3** which was isolated in 55% yield. The post-monoarylation of diphenyldithienylethene **3** by direct palladium catalyzed reaction was conducted at C2-atom of thiophene ring. The new compound **DTE-3-CHO-1** was synthesized with moderate yields (49%). The structures of **DTE-3-CHO-1** was confirmed by NMR and HR-MS spectroscopies.



Scheme 1. Synthetic strategy for **DTE-3-CHO-1**

4. Conclusions

We reported herein a new synthetic pathway for the preparation of diphenyldithienylethene. The direct palladium catalyzed C-H bond functionalization provided a convenient approach for regioselective monoraylation of diphenyldithienylethene. The structure of the obtained compound was confirmed by NMR and HR-MS spectral data. This method can be easily applied to prepare arylated diphenyldithienylethene with interesting photophysical properties. In addition, fluorescence and the aggregation induced emission study of this compound have shown promising results which will be published soon in due course.

❖ **Conflict of Interest:** Authors have no conflict of interest to declare.

❖ **Acknowledgement:** This research was funded by the Ministry of Education and Training under the grant number B2018-SPS-22.

REFERENCES

- Bottalico, D., Fiandanese, V., Marchese, G., & Punzi, A. (2009). Synthesis of Symmetrical Ketones from Grignard Reagents and 1,1'-Carbonyldiimidazole. *Synthesis*, 14, 2316–2318.
- Chang, Z. F., Jing, L. M., Liu, Y. Y., Liu, J. J., Ye, Y. C., Zhao, Y. S., Yuan, S.-C., & Wang, J. L. (2016). Constructing small molecular AIE luminophores through a 2, 2-(2, 2-diphenylethene-1, 1-diyl) dithiophene core and peripheral triphenylamine with applications in piezofluorochromism, optical waveguides, and explosive detection. *Journal of Materials Chemistry C*, 4, 8407–8415.
- Hong, Y., Lam, J. W. Y., & Tang, B. Z. (2011). Aggregation-Induced Emission. *Chemical Society Reviews*, 40, 5361–5388.
- Li, Y., Li, Z., Wang, Y., Compaan, A., Ren, T., & Dong, W.-J. (2013). Increasing the power output of a CdTe solar cell via luminescent down shifting molecules with intramolecular charge

- transfer and aggregation induced emission characteristics. *Energy & Environmental Science*, 6, 2907–2911.
- Lin, E. E., Wu, J.-Q., Schäfers, F., Su, X.-X., Wang, K.-F., Li, J.-L., Chen, Y., Zhao, X., Ti, H., Li, Q., Ou, T.-M., Glorius, F., & Wang, H. (2019). Regio- and stereoselective synthesis of tetra- and triarylethenes by *N*-methylimidodiacetyl boron-directed palladium-catalysed three-component coupling. *Communications Chemistry*, 2(34).
- Liu, X., Liang, G. (2017). Dual aggregation-induced emission for enhanced fluorescence sensing of furin activity *in vitro* and in living cells. *Chemical Communications*, 53, 1037–1040.
- Mei, J., Leung, N. L. C., Kwok, R. T. K., Lam, J. W. Y., & Tang, B. Z. (2015). Aggregation-Induced Emission: Together We Shine, United We Soar!. *Chemical Reviews*, 115(21), 11718–11940.
- Młostoń, G., Urbaniak, K., Pawlak, A., & Heimgartner, H. (2016). New Applications of Hetaryl Thioketones for the Synthesis of Hetaryl-Substituted Ethenes via 'Two-Fold Extrusion Reaction'. *Heterocycles*, 93, 127–139.
- Rananaware, A., Bhosale, R. S., Ohkubo, K., Patil, H., Jones, L. A., Jackson, S. L., Fukuzumi, S., Bhosale, S. V., & Bhosale, S. V. (2015). Tetraphenylethene-Based Star Shaped Porphyrins: Synthesis, Selfassembly, and Optical and Photophysical Study. *Journal of Organic Chemistry*, 80(8), 3832–3840.
- Shi, J., Deng, Q., Li, Y., Zheng, Z., Shangguan, H., Li, L., Huang, F., & Tang, B. (2019). Homogeneous probing of lipase and α -amylase simultaneously by AIEgens, *Chemical Communications*, 55, 6417–6420.
- Zhang, M., Yao, Y., Stang, P. J., & Zhao, W. (2020). Divergent and Stereoselective Synthesis of Tetraarylethenes from Vinylboronates. *Angewandte Chemie International Edition*, 59(45), 20090–20098.
- Zhang, G.-F., Wang, H., Aldred, M. P., Chen, T., Chen, Z.-Q., Meng, X., & Zhu, M.-Q. (2014). General Synthetic Approach toward Geminal-Substituted Tetraarylethene Fluorophores with Tunable Emission Properties: X-ray Crystallography, Aggregation-Induced Emission and Piezofluorochromism. *Chemistry of Materials*, 26, 4433–4446.
- Zhao, Z., Lu, P., Lam, J. W. Y., Wang, Z., Chan, C. Y. K., Sung, H. H. Y., Williams, I. D., Mab, Y., & Tang, B. Z. (2011). Molecular anchors in the solid state: Restriction of intramolecular rotation boosts emission efficiency of luminogen aggregates to unity. *Chemical Science*, 2, 672–675.
- Wang, X., Zhang, C., Xu, Y., He, Q., Mu, P., Chen, Y., Zeng, J., Wang, F., & Jiang, J. X. (2018). Conjugated Microporous Polytetra(2-Thienyl)ethylene as High Performance Anode Material for Lithium and Sodium-Ion Batteries. *Macromolecular Chemistry and Physics*, 219(7), 1700524–1700532.