

A METHOD OF APPROXIMATION FOR A ZERO OF MAXIMAL MONOTONE OPERATOR IN HILBERT SPACE

Phạm Thị Thu Hoài¹, Nguyễn Thị Thuy Hoa², Nguyễn Tất Thắng^{3*}

¹Vietnam Maritime University, ²Hanoi College of Home Affairs, ³Thai Nguyen University

ABSTRACT

In this paper, we introduce a new explicit iterative method for solving a variational inequality problem over the set of zeros for a maximal monotone operator in Hilbert space. By using two resolvents of the monotone operator at each iterate, we prove strong convergence of the method under a general condition on resolvent parameter.

Keywords: Maximal monotone operators; Nonexpansive mappings; Fixed points; Zero points; Variational inequalities

Received: 21/02/2020; **Revised:** 28/02/2020; **Published:** 29/02/2020

MỘT PHƯƠNG PHÁP XÁP XỈ CHO KHÔNG ĐIỂM CỦA TOÁN TỬ ĐƠN ĐIỀU CỰC ĐẠI TRONG KHÔNG GIAN HILBERT

Phạm Thị Thu Hoài¹, Nguyễn Thị Thuy Hoa², Nguyễn Tất Thắng³

¹Trường Đại học Hàng hải Việt Nam, ²Trường Đại học Nội vụ Hà Nội, ³Đại học Thái Nguyên

TÓM TẮT

Trong bài báo này chúng tôi đưa ra một phương pháp lặp hiện mới giải bài toán bất đẳng thức biến phân trên tập không điểm của toán tử đơn điệu cực đại trong không gian Hilbert. Bằng việc sử dụng hai toán tử giải của một toán tử đơn điệu tại mỗi bước lặp, chúng tôi chứng minh sự hội tụ mạnh của phương pháp dưới điều kiện suy rộng đặt lên tham số.

Từ khóa: Toán tử đơn điệu cực đại; ánh xạ không giãn; điểm bất động; không điểm; bất đẳng thức biến phân

Ngày nhận bài: 21/02/2020; **Ngày hoàn thiện:** 28/02/2020; **Ngày đăng:** 29/02/2020

* Corresponding author. Email: nguyentatthang.tnu@gmail.com

<https://doi.org/10.34238/tnu-jst.2020.02.2693>

1 Introduction

Let H be a real Hilbert space with inner product and norm denoted, respectively, by $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$. Let A be a maximal monotone operator in H . In this paper we assume that the set of zeros, $\Gamma := \{p \in \mathcal{D}(A) : 0 \in Ap\}$, is nonempty, where $\mathcal{D}(A)$ denotes the domain of A .

Finding a zero of a maximal monotone operator, i.e., finding a point

$$\text{finding a point } p \in \mathcal{D}(A) : \text{ such that } 0 \in Ap, \quad (1.1)$$

is an important part of the theory of monotone operators. A fundamental method for finding a zero point of a maximal monotone operator A in Hilbert space H , we can cite the proximal point one [1]. This method generates a sequence $\{x^k\}$ defined for each $k \geq 1$

$$x^{k+1} = J_k^A x^k + e^k \quad \text{or} \quad x^{k+1} = J_k^A(x^k + e^k) \quad (1.2)$$

where $x^1 \in H$, $J_k^A = (I + r_k A)^{-1}$, I is the identity mapping of H , $\{r_k\}$ is a sequence of real numbers such that $r_k \geq \varepsilon > 0$ for all $k \geq 1$ and e^k is an error vector, satisfying

$$\|x^{k+1} - J_k^A x^k\| \leq \varepsilon_k \quad \text{with} \quad \sum_{k=1}^{\infty} \varepsilon_k < \infty$$

or

$$\|x^{k+1} - J_k^A x^k\| \leq \eta_k \|x^{k+1} - x^k\| \quad \text{with} \quad \sum_{k=1}^{\infty} \eta_k < \infty. \quad (1.3)$$

Methods (1.2)-(1.3) converge only weakly to a zero of A in infinite-dimensional Hilbert spaces, in general (see, [2]). In order to have a strong convergence sequence $\{x^k\}$ from the method, several modifications of (1.2) were proposed in [3–5]. Kamimura and Takahashi [3] introduced a method, in there two sequences $\{x^k\}$ and $\{y^k\}$ are built from a starting point x^1 as follows:

$$y^k \approx J_k^A x^k, \quad \|y^k - J_k^A x^k\| \leq \delta_k, \quad x^{k+1} = t_k u + (1 - t_k) y^k, \quad k \geq 1, \quad (1.4)$$

where u is a fixed point in H . The sequence $\{x^k\}$ so generated, as $k \rightarrow \infty$, is strongly convergent to $P_\Gamma u$, the metric projection of u on the set Γ , under the following conditions:

(C1) $t_k \in (0, 1)$ for all $k \geq 1$, $\lim_{k \rightarrow \infty} t_k = 0$ and $\sum_{k=1}^{\infty} t_k = \infty$;

(C2) $r_k \in (0, \infty)$ for all $k \geq 1$ and $\lim_{k \rightarrow \infty} r_k = \infty$; and

(C3) $\sum_{k=1}^{\infty} \delta_k < \infty$.

Xu [5] extended the prox-Tikhonov method of Lehdili and Moudafi [4] in the following way

$$x^{k+1} = J_k^A(t_k u + (1 - t_k)x^k + e^k). \quad (1.5)$$

Further, Boikanyo and Morosanu [6] showed that (1.5) is equivalent to

$$y^{k+1} = t_k u + (1 - t_k)J_k^A y^k + e^k, \quad (1.6)$$

and proved a strong convergence result for $\{y^k\}$, defined by (1.6), to $P_\Gamma u$, if there hold conditions (C1), (C2) and

(C3') either (C3) with $\delta_k = \|e^k\|$ or $\lim_{k \rightarrow \infty} (\|e_k\|/t_k) = 0$.

In this paper, we introduce a new modifications of (1.2),

$$x^{k+1} = J_k^A J_c^A(t'_k u + (1 - t'_k)x^k) + e^k \quad (1.7)$$

where $J_c^A = (I + cA)^{-1}$ and c is any fixed positive real number. We will show that method (1.7) is particular case of the following method,

$$z^{k+1} = J_k^A J_c^A(I - t_k \mu F)z^k + e^k, \quad (1.8)$$

proposed to solve a problem of finding a point

$$p_* \in \Gamma \quad \text{such that} \quad \langle Fp_*, p_* - p \rangle \leq 0 \quad \forall p \in \Gamma, \quad (1.9)$$

where $\mu \in (0, 2\eta/L^2)$ is a constant and $F : H \rightarrow H$ is an η -strongly monotone and L -Lipschitz continuous operator with $\eta, L > 0$.

The paper is organized as follows. In Section 2, we list some related facts that will be used in our result. In Section 3, we prove strong convergence of our main method and show that their special case is new contraction and generalized proximal point method, that converge strongly to a zero under a general condition on the resolvent parameter.

2 Preliminaries

In this section, we introduce some mathematical symbols, definitions, and lemmas which can be used in the proof of our main result.

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|$. In what follows, we write $x^k \rightharpoonup x$ to indicate that the sequence $\{x^k\}$ converges weakly to x while $x^k \rightarrow x$ indicate that the sequence $\{x^k\}$ converges strongly to x .

First, we know that, for any Hilbert space H ,

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle \quad \forall x, y \in H.$$

Let C be a nonempty, closed and convex subset of H . We know that, for each $x \in H$, there is a unique $P_C x \in C$ such that

$$\|x - P_C x\| = \inf_{u \in C} \|x - u\|, \quad (2.1)$$

and the mapping $P_C : H \rightarrow C$ defined by (2.1) is called the metric projection from H onto C . Moreover, we have

$$\langle x - P_C x, y - P_C x \rangle \leq 0, \quad \forall x \in H, y \in C, \quad (2.2)$$

(see, for example, [7, Section 3]).

Let $F : H \rightarrow H$ be a mapping. F is said to be L -Lipschitz continuous and η -strongly monotone when the following conditions are satisfied:

$$\|Fx - Fy\| \leq L\|x - y\| \quad \text{and} \quad \langle Fx - Fy, x - y \rangle \geq \eta\|x - y\|^2$$

for all $x, y \in H$, where L and η are some positive constants. F is said to be contraction operator, if $0 \leq L < 1$ and nonexpansive, if $L = 1$.

Lemma 2.1 [see, [8]] Let H be a real Hilbert space and let F be an η -strongly monotone and L -Lipschitz continuous operator on H with some positive constants η and L . Then, for a fixed number $\mu \in (0, 2\eta/L^2)$ and any $t \in (0, 1)$, $I - t\mu F$ is a contraction with contractive constant $1 - t\tau$, where $\tau = 1 - \sqrt{1 - \mu(2\eta - \mu L^2)}$.

We introduce some definitions and propositions about set-valued mappings. Let A be a set-valued operator of H into 2^H with domain $\mathcal{D}(A) = \{x \in H : A(x) \neq \emptyset\}$, range $\mathcal{R}(A) = \bigcup_{x \in \mathcal{D}(A)} Ax$, and the inverse of A is $A^{-1}(y) = \{x \in H : y \in A(x)\}$. $A : H \rightarrow 2^H$ is said to be

- (i) monotone if $\langle u - v, x - y \rangle \geq 0 \quad \forall x, y \in \mathcal{D}(A), u \in A(x), v \in A(y)$;
- (ii) maximal monotone if it is monotone and the graph

$$\mathcal{G}(A) = \{(x, y) \in H \times H : x \in \mathcal{D}(A), y \in A(x)\}$$

of A is not properly contained in the graph of any other monotone operator on $\mathcal{D}(A)$.

For a monotone mapping A , we define its resolvent J_r^A by

$$J_r^A := (I + rA)^{-1} : \mathcal{R}(I + rA) \subset H \rightarrow \mathcal{D}(A),$$

where $r > 0$, I is the identity operator on H .

A fixed point of the mapping $F : C \rightarrow C$ is a point $x \in C$ such that $Fx = x$. The set of all fixed points of the mapping F is denoted by $\text{Fix}(F)$.

Lemma 2.2 [9, Section 7] Let H be a real Hilbert space. If $A : H \rightarrow H$ is a maximal monotone operator,

(i) J_r^A is nonexpansive, single-valued mapping and $\text{Fix}(J_r^A) = A^{-1}(0)$ for each $r > 0$, and

(ii)

$$\|J_r^A x - p\|^2 \leq \|x - p\|^2 - \|J_r^A x - x\|^2 \quad \forall x \in H, p \in A^{-1}(0).$$

Lemma 2.3 [see, [10]] Let $\{a_k\}$, $\{b_k\}$ and $\{c_k\}$ be sequences of real numbers such that, for all $k \geq 1$, $a_{k+1} \leq (1 - b_k)a_k + b_k c_k$; $a_k \geq 0$;

(i) b_k satisfies a condition of type **(T)**; and

(ii) either $\sum_{k=1}^{\infty} b_k |c_k| < \infty$ or $\limsup_{k \rightarrow \infty} c_k \leq 0$.

Then, $\lim_{k \rightarrow \infty} a_k = 0$.

Lemma 2.4 [see, [11]] Let $\{a_k\}$ be a sequence of real numbers with a subsequence $\{k_l\}$ of $\{k\}$ such that $a_{k_l} < a_{k_l+1}$ for all $l \in N_+$. Then, there exists a nondecreasing sequence $\{m_k\} \subseteq N_+$ such that $m_k \rightarrow \infty$, $a_{m_k} \leq a_{m_k+1}$ and $a_k \leq a_{m_k+1}$ for all (sufficiently large) numbers $k \in N_+$. In fact, $m_k = \max\{l \leq k : a_l \leq a_{l+1}\}$.

Proposition 2.1 [see, [12, 13]] Let H and F be as in Lemma 2.1 and let T be a nonexpansive operator on H such that $\text{Fix}(T) \neq \emptyset$. Then, for any bounded sequence $\{x^k\} \subset H$ such that $\lim_{k \rightarrow \infty} \|Tx^k - x^k\| = 0$, we have

$$\limsup_{k \rightarrow \infty} \langle Fp_*, p_* - x^k \rangle \leq 0, \quad (2.3)$$

where $p_* \in \text{Fix}(T)$, solving (1.9) with Γ replaced by $\text{Fix}(T)$.

3 Main Results

First of all, we have the following results.

Theorem 3.1 Let A be a maximal monotone operator in a real Hilbert space H such that $\Gamma := \{p \in \mathcal{D}(A) : 0 \in Ap\} \neq \emptyset$ and let F with μ be as in Lemma 2.1. Assume that there hold conditions (C1), (C3'), and

(C2') $\{r_k\}$ is any sequence of numbers in $(0, \infty)$.

Then, the sequence $\{z^k\}$, defined by (1.8), as $k \rightarrow \infty$, converges strongly to the unique solution p_* , solving (1.9).

Proof We consider an exact variant of (1.8), that is,

$$x^{k+1} = J_k^A J_c^A (I - t_k \mu F) x^k. \quad (3.1)$$

Clearly, from (1.8), (3.1), the nonexpansive property of J_k^A and Lemma 2.1, we get the following inequality:

$$\begin{aligned} \|z^{k+1} - x^{k+1}\| &= \|J_k^A J_c^A (I - t_k \mu F) z^k + e^k - J_k^A J_c^A (I - t_k \mu F) x^k\| \\ &\leq (1 - t_k \tau) \|z^k - x^k\| + \|e^k\|. \end{aligned}$$

According to conditions (C1) and (C3'), we apply Lemma 2.2 to conclude that $\|z^k - x^k\| \rightarrow 0$ as $k \rightarrow \infty$. Hence, to show the desired result, it suffices to prove that $\{x^k\}$ converges strongly to p_* as $k \rightarrow \infty$. For this purpose, we first prove that $\{x^k\}$ is bounded. Indeed, for a fixed point $p \in \Gamma$, by Lemma 2.1, we have

$$\begin{aligned} \|x^{k+1} - p\| &= \|J_k^A J_c^A (I - t_k \mu F) x^k - J_k^A J_c^A p\| \leq (1 - t_k \tau) \|x^k - p\| + t_k \mu \|Fp\| \\ &\leq \max \{\|x^1 - p\|, \mu \|Fp\| / \tau\}, \end{aligned}$$

by mathematical induction. Therefore, the sequence $\{x^k\}$ is bounded, and so are the sequences $\{Fx^k\}$ and $\{y^k\}$ where $y^k := (I - t_k \mu F)x^k$. Without loss of generality, we can assume that they are bounded by a positive constant M_1 .

Further, we estimate the value $\|x^{k+1} - p\|^2$ as follows.

$$\begin{aligned} \|x^{k+1} - p\|^2 &= \|J_k^A J_c^A y^k - J_k^A p\|^2 \leq \|J_c^A y^k - p\|^2 \leq \|y^k - p\|^2 - \|J_c^A y^k - y^k\|^2 \\ &= \|(I - t_k \mu F)x^k - p\|^2 - \|J_c^A y^k - y^k\|^2 \leq (1 - t_k \tau) \|x^k - p\|^2 + 2t_k \mu \langle Fp, p - y^k \rangle - \|J_c^A y^k - y^k\|^2. \end{aligned} \quad (3.2)$$

We need only consider two cases.

Case 1. There exists an integer $k_0 \geq 1$ such that $\|x^{k+1} - p\| \leq \|x^k - p\|$ for all $k \geq k_0$.

Then, $\lim_{k \rightarrow \infty} \|x^k - p\|$ exists. From (3.2), we can write that

$$\|J_c^A y^k - y^k\|^2 \leq \|x^k - p\|^2 - \|x^{k+1} - p\|^2 + 2t_k \mu M \quad (3.3)$$

where $M \geq \|Fp\|(\|p\| + M_1)$. Since $\lim_{k \rightarrow \infty} \|x^k - p\|$ exists and $t_k \rightarrow 0$, letting k tend to infinity in (3.3), we get that $\lim_{k \rightarrow \infty} \|J_c^A y^k - y^k\| = 0$. This together with $\|y^k - x^k\| \leq t_k \mu M_1$ implies

that $\lim_{k \rightarrow \infty} \|J_c^A x^k - x^k\| = 0$. By using Proposition 2.1 with $T = J_c^A$, we obtain inequality (2.3). Now, from (3.2) with $p = p_*$, we know that

$$\begin{aligned} \|x^{k+1} - p_*\|^2 &\leq (1 - t_k \tau) \|x^k - p_*\|^2 + 2t_k \mu \langle Fp_*, p_* - x^k + t_k \mu Fx^k \rangle \\ &= (1 - b_k) \|x^k - p_*\|^2 + b_k c_k \quad \text{for all } k \geq k_0, \end{aligned}$$

where $b_k = t_k \tau$ and $c_k = \frac{2\mu}{\tau} [\langle Fp_*, p_* - x^k \rangle + t_k \mu \|Fp_*\| M_1]$, from which and Lemma 2.2 we obtain that $\|x^k - p_*\| \rightarrow 0$.

Case 2. There exists a subsequence $\{k_l\}$ of $\{k\}$ such that $\|x^{k_l} - p\| < \|x^{k_l+1} - p\|$ for all $l \in N_+$. Hence, by Lemma 2.4, there exists a nondecreasing sequence $\{m_k\} \subseteq N_+$ such that $m_k \rightarrow \infty$,

$$\|x^{m_k} - p\| \leq \|x^{m_k+1} - p\| \quad \text{and} \quad \|x^k - p\| \leq \|x^{m_k+1} - p\| \quad (3.4)$$

for each $k \in N_+$. Then, from (3.2) and the first inequality in (3.4), we know that

$$\|x^{m_k} - p\|^2 \leq \frac{2\mu}{\tau} \langle Fp, p - y^{m_k} \rangle. \quad (3.5)$$

On the other hand, again from (3.2) the first inequality in (3.4), we have also that

$$\|J_c^A y^{m_k} - y^{m_k}\|^2 \leq 2t_{m_k} \mu M.$$

Therefore, $\lim_{k \rightarrow \infty} \|J_c^A y^{m_k} - y^{m_k}\| = 0$, and hence, by Proposition 2.1,

$$\limsup_{k \rightarrow \infty} \langle Fp_*, p_* - y^{m_k} \rangle \leq 0,$$

from which and (3.5) with p replaced by p_* , it follows that

$$\lim_{k \rightarrow \infty} \|x^{m_k} - p_*\| = 0. \quad (3.6)$$

Finally, from (3.2) with k and p replaced, respectively, by m_k and p_* , we can write that

$$\|x^{m_k+1} - p_*\|^2 \leq (1 - t_{m_k} \tau) \|x^{m_k} - p_*\|^2 + 2t_{m_k} \mu \langle Fp_*, p_* - y^{m_k} \rangle.$$

By virtue of (3.6) and $t_{m_k} \rightarrow 0$, $\lim_{m \rightarrow \infty} \|x^{m_k+1} - p_*\|^2 = 0$, which together with the second inequality in (3.4) implies that $\lim_{k \rightarrow \infty} \|x^k - p_*\| = 0$. This completes the proof.

Remark 3.1 Clearly, the mapping $F = I - f$, where $f = \tilde{a}I + (1 - \tilde{a})u$ with a fixed number $\tilde{a} \in (0, 1)$ and a fixed point $u \in H$, is an η -strongly monotone and L -Lipschitz continuous operator with $\eta = 1 - \tilde{a}$ and $L = 1 + \tilde{a}$. Then, replacing F in (1.8) by $I - f$ and re-denoting $z^k := x^k$ with $t'_k := t_k \mu (1 - \tilde{a})$, we obtain (1.7). Now, putting $y^k = (I - t_k \mu F)x^k$ in (1.8), we get that

$$y^{k+1} = (I - t_{k+1} \mu F)x^{k+1} = (I - t_{k+1} \mu F)(J_k^A J_c^A y^k + e^k).$$

4 Conclusion

We have presented an iterative method for finding a point in the zero set of a maximal monotone mapping in Hilbert space, that solves a variational inequality problem, involving an η -strongly monotone and L -Lipschitz continuous operator on H for some positive constants η and L . As consequences, new generalized and contraction proximal point algorithm with a any sequence of positive numbers for the resolvent parameter have been obtained.

REFERENCES

- [1]. R.T. Rockafellar, "Monotone operators and the proximal point algorithm", *SIAM J. Control Optim.*, **14**(5), pp. 877–898, 1976.
 - [2]. O. Guler, "On the convergence of the proximal point algorithm for convex minimization", *SIAM J. Control Optim.*, **29**(2), pp. 403–419, 1991.
 - [3]. S. Kakimura, W. Takahashi, "Approximating solutions of maximal monotone operators in Hilbert spaces", *J. Approx. Theory*, **106**(2), pp. 226–240, 2000.
 - [4]. N. Lehdili, A. Moudafi, "Combining the proximal point algorithm and Tikhonov regularization", *Optimization*, **37**(3), pp. 239–252, 1996.
 - [5]. H.K. Xu, "A regularization method for the proximal point algorithm", *J. Glob. Optim.*, **36**(1), pp. 115–125, 2006.
 - [6]. O.A. Boikanyo, G. Morosanu, "A proximal point algorithm converging strongly for general errors", *Optim. Lett.*, **4**(4), pp. 635–641, 2010.
 - [7]. K. Goebel, S. Reich, *Uniform Convexity, Hyperbolic Geometry, and Nonexpansive Mappings*, Marcel Dekker, New York, 1984.
 - [8]. I. Yamada, "The hybrid steepest descent method for the variational inequality problem over the intersection of fixed point sets of nonexpansive mappings" in *Inherently Parallel Algorithms in Feasibility and Optimization and Their Applications* (D.Butnariu, Y. Censor and S. Reich, Eds). North-Holland, Amsterdam, pp. 473–504, 2001.
 - [9]. S. Reich, "Extension problems for accretive sets in Banach spaces", *J. Functional Anal.*, **26**, pp. 378–395, 1977.
 - [10]. H.K. Xu, "Iterative algorithms for nonlinear operators", *J. Lond. Math. Soc.*, **66**(1), pp. 240–256, 2002.
-

-
- [11]. P.E. Maingé, "Strong convergence of projected subgradient methods for nonsmooth and nonstrictly convex minimization", *Set-Valued Var. Anal.*, **16**(7-8), pp. 899–912, 2008.
- [12]. N. Buong, V.X. Quynh, N.T.T. Thuy, "A steepest-descent Krasnosel'skii–Mann algorithm for a class of variational inequalities in Banach spaces", *J. Fixed Point Theory and Appl.*, **18**(3), pp. 519–532, 2016.
- [13]. N. Buong, N.S. Ha, N.T.T. Thuy, "A new explicit iteration method for a class of variational inequalities", *Numer. Algor.*, **72**(2), pp. 467–481, 2016.
-